

Observational Versus Randomized:

Kotecha, Dipak; Altman, Douglas G; Rosano, Giuseppe; Flather, Marcus D

DOI:

[10.1016/j.jchf.2017.02.013](https://doi.org/10.1016/j.jchf.2017.02.013)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Kotecha, D, Altman, DG, Rosano, G & Flather, MD 2017, 'Observational Versus Randomized: Sliding Toward Nonevidence-Based Medicine', *JACC: Heart Failure*, vol. 5, no. 5, pp. 395-396.
<https://doi.org/10.1016/j.jchf.2017.02.013>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Letter in response to **Cadrin-Tourigny et al.**, Decreased Mortality With Beta-Blockers in Patients With Heart Failure and Coexisting Atrial Fibrillation: An AF-CHF Substudy; **JACC Heart Fail** 2017;10.1016/j.jchf.2016.10.015

Title: Observational versus randomized - Sliding towards non-evidence based medicine

Authors: Dipak Kotecha MD PhD^{1,2}, Douglas G Altman DSc³, Giuseppe Rosano MD PhD^{4,5} and Marcus D Flather MD⁶

From the: (1) University of Birmingham Institute of Cardiovascular Sciences, Birmingham, UK; (2) Centre of Cardiovascular Research and Education in Therapeutics, Monash University, Melbourne, Australia; (3) Centre for Statistics in Medicine, Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, Oxford, UK; (4) Department of Medical Sciences, IRCCS San Raffaele Pisana, Roma, Italy; (5) Cardiovascular and Cell Science Institute, St George's University of London, UK; (6) Norwich Medical School, University of East Anglia, Norwich, UK.

Word count: 497 (max. 500)

Corresponding author: Dipak Kotecha MD PhD MRCP PCAP FESC FHEA

University of Birmingham Institute of Cardiovascular Sciences, Medical School, Vincent Drive,
Birmingham, B15 2TT, UK

d.kotecha@bham.ac.uk; +44 121 371 8122

Conflicts: DK is the Chief Investigator for the RAte control Therapy Evaluation in permanent Atrial Fibrillation trial (RATE-AF; [NCT02391337](#)), the lead for the Beta-blockers in Heart Failure Collaborative Group (BB-meta-HF; [NCT00832442](#)), and has received research grants from Menarini, professional development support from Daiichi-Sankyo and lecture fees from AtriCure. MDF reports personal fees from AstraZeneca and grants from Novartis, all outside the submitted work. DGA and GR have nothing to disclose.

We welcome the addition of further data in the field of atrial fibrillation (AF) and heart failure (HF), a combination of conditions that results in difficult management decisions and worse outcomes, both for patients with reduced (1) and preserved left ventricular ejection fraction LVEF.(2) However, it remains important to adhere to some fundamental aspects of evidence-based medicine. Recent years have seen an abundance of subgroup analyses attempting to answer questions relating to treatment effects in non-randomized studies. These analyses, often using propensity matching or other statistical adjustment, have become commonplace and are often mistakenly considered to be as important as randomized controlled trials (RCTs). Regardless of analysis method, observational data should only be used to generate hypotheses about treatment effects (3,4), and we should resist the temptation to analyze datasets simply because of availability. Ignoring the weaknesses of such studies, and incorrect interpretation, can subsequently lead to confounded conclusions.

In the case of the article by Cadrin-Tourigny et al.(5), the patients were not randomized to beta-blockers and hence there is confounding at both the patient and physician level that hampers external validity. Doctors typically give beta-blockers to patients at lower risk, confirmed in this study as being younger in age, with more non-ischemic cardiomyopathy, less time in AF, and higher use of anticoagulation and defibrillators, all factors associated with lower mortality. Whilst propensity-matched analysis can be useful to mitigate minor differences in demographics, it was not designed to account for different patient populations, or for exposures that interact with the outcome. This has been demonstrated for digoxin therapy (the inverse of beta-blockers, in which clinicians tend to prescribe to higher risk patients), where propensity-matching was unable to replicate the results of RCTs.(6) Confounding may also explain why the authors of this paper found such discrepant findings for death and hospitalization. Further, only 57% of their population were actually in AF at

the time of analysis. We have already shown how effective beta-blockers are in preventing AF (and therefore subsequent adverse outcomes) for HF patients in sinus rhythm.(7) A few methodological issues also arise on detailed review, including the divergence in matched groups, omission of paroxysmal versus persistent AF from the standardized difference plot (with clearly more than 10% difference), and misrepresentation in the abstract about the sample size (n=655 not 1376, with just 95 deaths without beta-blockers and 136 deaths on beta-blockers).

We considered the requirement for AF on the baseline ECG a strength of our previous analysis, which demonstrated a significant interaction in beta-blocker efficacy according to heart rhythm using data from double-blind placebo-controlled RCTs in patients with HF and reduced LVEF.(7) Using systematic, carefully checked and harmonized individual patient data from ten trials, we identified no significant benefit from beta-blockers in over 3000 patients with concomitant AF, consistent across all outcomes studied and based on a published design paper and pre-specified analysis plan.

Where, as a body of clinical scientists, do we go from here? The answer lies in new RCTs, rather than more analyses of observational data.

References

1. Kotecha D, Piccini JP. Atrial fibrillation in heart failure: what should we do? *Eur Heart J* 2015;36:3250-7.
2. Kotecha D, Lam CS, Van Veldhuisen DJ, Van Gelder IC, Voors AA, Rienstra M. Heart Failure With Preserved Ejection Fraction and Atrial Fibrillation: Vicious Twins. *J Am Coll Cardiol* 2016;68:2217-2228.
3. Vandenbroucke JP. Observational research, randomised trials, and two views of medical science. *PLoS Med* 2008;5:e67.
4. Greenland S. Randomization, statistics, and causal inference. *Epidemiology* 1990;1:421-9.
5. Cadrin-Tourigny J, Shohoudi A, Roy D et al. Decreased Mortality With Beta-Blockers in Patients With Heart Failure and Coexisting Atrial Fibrillation: An AF-CHF Substudy. *JACC Heart Fail* 2017;10.1016/j.jchf.2016.10.015.
6. Ziff OJ, Lane DA, Samra M et al. Safety and efficacy of digoxin: systematic review and meta-analysis of observational and controlled trial data. *BMJ* 2015;351:h4451.
7. Kotecha D, Holmes J, Krum H et al. Efficacy of beta blockers in patients with heart failure plus atrial fibrillation: an individual-patient data meta-analysis. *Lancet* 2014;384:2235-43.